

****FULL TITLE****

*ASP Conference Series, Vol. **VOLUME**, **YEAR OF PUBLICATION***

****NAMES OF EDITORS****

Composition of the galactic center star cluster

Rainer M. Buchholz¹, Rainer Schödel², Andreas Eckart^{1,3}

Abstract. We present a population analysis of the nuclear stellar cluster of the Milky Way based on adaptive optics narrow band spectral energy distributions. We find strong evidence for the lack of a stellar cusp and a similarity of the late type luminosity function to the bulge KLF.

1. Observations and classification

We observed the central parsec of the Galactic center using an H band filter and seven intermediate band filters covering the K band with the AO assisted instrument NACO at the ESO VLT on Paranal. This yielded eight point spectral energy distributions (SEDs) for 5914 sources. The spectral feature we used for stellar classification is the CO bandhead absorption ($\lambda > 2.24\mu\text{m}$), which allows the separation of late and early type stars. We determined its presence by fitting the SEDs with an extinguished blackbody. Our method produces reliable results for stars as faint as 15.5 mag in the K band (K2III or B2V stars, considering extinction and distance modulus), much deeper than previous studies. We classified 322 stars as early type candidates and 2955 as late type.

2. Results

For the first time, it is now possible to study the early and late type population separately at this depth. Only the latter can be expected to be dynamically relaxed due to its age (Gyrs). Our new results show that the late type population indeed lacks the predicted cusp structure, showing a core instead (see e.g. Merritt 2009). The steep decline of the projected early type density towards the outer parts of the cluster (e.g. Genzel et al. 2003; Paumard et al. 2006) could be confirmed here as well (for both see Fig.1). The total K band luminosity function of the central parsec is considerably flatter than that of the galactic bulge, with a power law index $\alpha_{GC} = 0.23 \pm 0.02$ (Schödel et al. 2007) compared to $\alpha_{bulge} = 0.3$ (Figer et al. 2004). The power law slope we fitted to the late type KLF alone agrees very well with that of the bulge, a surprising result since the GC cluster is assumed to be a dynamically separate entity (see Fig.1).

¹I. Physikalisches Institut, Universität zu Köln, Zöllicher Str. 77, 50937 Köln, Germany, email: buchholz,eckart@ph1.uni-koeln.de

²Instituto de Astrofísica de Andalucía (IAA)-CSIC, Glorieta de la Astronomía s/n, E-18008 Granada, Spain, email: rainer@iaa.es

³Max-Planck-Institut für Radioastronomie, Auf dem Hügel 69, 53121 Bonn, Germany




Figure 1. Left: Projected densities of different classes of sources, green: early type stars, fitted with $\beta = -1.49 \pm 0.12$, resp. $\beta_1 = -1.1 \pm 0.1$ and $\beta_2 = -3.5 \pm 0.6$ for $R_1 < 10'' < R_2$, red: late type stars, fitted with $\beta_1 = 0.17 \pm 0.09$ and $\beta_2 = -0.70 \pm 0.09$ for $R_1 < 6'' < R_2$, blue: all identified stars, light blue: Paumard et al. (2006) sources, Right: KLFs for different populations in the GC, with fitted power laws.

We can also confirm the much flatter KLF found for the early type stars (e.g. Paumard et al. 2006) to a much greater depth ($K < 15.5$ instead of $K < 13-14$).

3. Conclusions

These new results have enhanced our knowledge of the stellar population in the GC considerably, although several points remain unclear. Instead of a cusp, we find a depletion of giants in the very center, as already found by e.g. Haller et al. (1996), for giants with $K < 12$, with Do et al. (2009); Bartko et al. (2010) confirming our own results spectroscopically only recently. Several possible causes for this have been suggested, such as collisions/envelope stripping, mass segregation and IMBH/SMBH infall scenarios (see e.g. Dale et al. 2009). The fact that the power law slopes of the bulge and the late type KLF agree within errors points to a similar mass function and stellar evolution. How this agrees with both being dynamically separate entities needs to be determined. For our further results and a more detailed treatment of the issues mentioned above, please see Buchholz et al. (2009).

References

- Bartko, H., Martins, F., Trippe, S., Fritz, T., Genzel, R., et al. 2010, ApJ, 708, 834B
- Buchholz, R. M., Schödel, R., Eckart, A. 2009, A&A, 499, 483
- Dale, J., Davies, M., Church, R., Freitag, M. 2009, MNRAS 393, 1016
- Do, T., Ghez, A., Morris, M., Lu, J., et al. 2009, ApJ 703, 1323D
- Figer, D., Rich, R., Sunsoo, S., Morris, M., & Serabyn, E. 2004, ApJ, 601, 319
- Genzel, R., Schödel, R., Ott, T., Eisenhauer, F., et al. 2003, ApJ, 594, 812
- Haller, J. W., Rieke, M. J., Rieke, G. H., et al. 1996, ApJ, 456, 194
- Merritt, D. 2009, arXiv:0909.1318
- Paumard, T., Genzel, R., Martins, F., et al. 2006, ApJ, 643, 1011-1035
- Schödel, R., Eckart, A., Alexander, T., et al. 2007, A&A, 469, pp.125-146